Cliente: RUBIÓ, S. A.	Impresión:
Especialidad: CHOLEST USP LIGHT	Black / Cyan / Magenta / Yellow / Reserva Barniz / Troquel
Ref. cliente: 38026005va0120	
Sust. a.: -	
Tamaño: 100 x 130 x 145	
Troquel: BCN0243C.cf2	
Modificación: Nueva	OBSERVACIÓN: Tener en cuenta los valores para impresión
	Fecha: 13/01/2021 1

PANS

(02)00000380260056(99)0120

Preparation of Cholestyramine for Oral Suspension USP, Light
Cholestyramine for Oral Suspension USP, Light can be easily mixed with highly fluid foods or fruits.

1. Pour contents of one Cholestyramine for Oral Suspension USP, Light pouch in a bowl.
2. Add at least 6 ounces of applesauce or other food.
3. Mix well.
4. The slightly-textured Cholestyramine for Oral Suspension USP, Light mixture is now ready to eat.

NOTE: More than one pouch can be mixed at one time depending upon the amount of food you wish to consume when taking your Cholestyramine for Oral Suspension USP, Light. You may mix to suit your individual taste.

0120

60 Single-Dose Pouches NDC 51224-009-20

Cholestyramine for Oral Suspension USP, Light

Powder

4 grams cholestyramine resin USP, per pouch*

*each pouch contains 4 grams cholestyramine resin USP

Orange Flavor
Rx Only

Usual Dosage: See accompanying literature.
WARNING: This package is not child-resistant. **KEEP OUT OF THE REACH OF CHILDREN.**

tagi PHARMA

Healthful Hints

- Maintain diet and exercise as directed by your physician.
- Find a suitable time to take your medication daily (it may be useful to take Cholestyramine for Oral Suspension USP, Light with meals).
- Cholestyramine for Oral Suspension USP, Light can be mixed and refrigerated for three days (remember to mix well each time before drinking or eating).
- Cholestyramine for Oral Suspension USP, Light can also be easily mixed in a blender.
- Ask your pharmacist or physician for new and different ways to prepare Cholestyramine for Oral Suspension USP, Light.

Contains the artificial sweetener Aspartame.
WARNINGS PHENYLKETONURICS: CONTAINS PHENYLALANINE 16.8 mg per 4.8 g DOSE.

Preparation of Cholestyramine for Oral Suspension USP, Light
Cholestyramine for Oral Suspension USP, Light can be mixed with your choice of noncarbonated beverage.

1. Pour contents of one Cholestyramine for Oral Suspension USP, Light pouch into a glass or cup.
2. Add 2 ounces of your favorite noncarbonated beverage (orange juice is a popular choice) and stir vigorously.
3. Add at least 2 to 4 more ounces of noncarbonated beverage to suit individual taste and stir vigorously again.
4. The slightly-textured Cholestyramine for Oral Suspension USP, Light mixture is now ready to drink.

NOTE: More than one pouch can be mixed at one time depending upon the amount of noncarbonated beverage you wish to consume when taking your Cholestyramine for Oral Suspension USP, Light. You may mix to suit your individual taste.

38026005

60 Single-Dose Pouches NDC 51224-009-20
Cholestyramine for Oral Suspension USP, Light
Powder

4 grams cholestyramine resin USP, per pouch*
*each pouch contains 4 grams cholestyramine resin USP
Rx only

Store at 20°C to 25°C (68° to 77°F) [see USP Controlled Room Temperature].
Always mix Cholestyramine for Oral Suspension USP, Light with water, or the noncarbonated beverage of your choice, or other highly fluid foods or fruits before using.

Manufactured by
Laboratorios Rubió, S.A.
Industria, 29. Pol. Ind. Comte de Sert
08755 Castellbisbal (Barcelona)
Spain

Manufactured for
TAGI Pharma, Inc.
South Beloit, IL 61080

Rev. 12/20

3 51224 00920 2

1/13/21
1/13/21
1/13/21